

AD-A103 039

WASHINGTON UNIV SEATTLE DEPT OF CHEMISTRY F/G 7/3
PYROLYSIS OF 1-IODOPROPANE BY THE VARIABLE ENCOUNTER METHOD.(U)
AUG 81 F C WOLTERS, K J CHAO, B S RABINOVITCH N00014-75-C-0690
TR-21 NL

UNCLASSIFIED

1 of 1
AD
AD 700 039



END
DATE
FILMED
9-81
DTIC

LEVEL

12

Pyrolysis of 1-Iodopropane by the Variable Encounter Method

F. C. Wolters, K.-J. Chao, and B. S. Rabinovitch

Department of Chemistry BG-10
University of Washington
Seattle, WA 98195

Technical Report No. NR092-549-TR21
Contract N00014-75-C-0690, NR-092-549

August 1, 1981

Prepared for Publication in the
International Journal of Chemical Kinetics



OFFICE OF NAVAL RESEARCH
Department of the Navy
Code 473
800 N. Quincy
Arlington, VA 22217

Reproduction in whole or in part is permitted for any purpose of
the United State Government. This document has been approved for
public release; its distribution is unlimited.

81 8 18 189

AD A103039

DTIC FILE COPY



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER NR092-549-TR21	2. GOVT ACCESSION NO. AD-A103	3. RECIPIENT'S CATALOG NUMBER 039	
4. TITLE (and Subtitle) Pyrolysis of 1-Iodopropane by the Variable Encounter Method		5. TYPE OF REPORT & PERIOD COVERED Technical	
		6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) F. C. Wolters, K.-J. Chao, and B. S. Rabinovitch		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0690 NR 092-549	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Professor B. S. Rabinovitch Department of Chemistry BG-10 University of Washington Seattle, WA 98195		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 743 Department of the Navy 800 N. Quincy Arlington, VA 22217		12. REPORT DATE August 1, 1981	
		13. NUMBER OF PAGES 74	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; its distribution is unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES Prepared for publication in Intern. J. Chem. Kinetics			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Accommodation Polarmolecule Collisional Single Collision Energy Transfer Surfaces Gases Unimolecular Reaction Iodopropane Vibrational Relaxation			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The pyrolysis of 1-iodopropane has been studied by the Variable Encounter Method (VEM) at temperatures from 625 K to 840 K. Deactivating wall colli- sions are found to be stronger for this molecule than for the non-polar hydrocarbons previously studied in the same temperature range. Energy trans- fer collision efficiency declines with rise of wall temperature. The results of this study are compared with earlier steady-state work.			

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
(N 0102 (1-014) 6001)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Pyrolysis of 1-Iodopropane by the Variable Encounter Method*

F. C. Wolters,[†] K.-J. Chao[‡] and B. S. Rabinovitch

Department of Chemistry BG-10
University of Washington, Seattle, Washington 98195

Abstract

The pyrolysis of 1-iodopropane has been studied by the Variable Encounter Method (VEM) at temperatures from 625 K to 840 K. Deactivating wall collisions are found to be stronger for this molecule than for the hydrocarbons previously studied in this temperature range. The results of this study are compared with earlier steady-state work.

Accession For	
NTIS	<input checked="" type="checkbox"/>
NTIC	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Director	
Approved	
A	


* Work supported by the Office of Naval Research

[†] Present address: Clorox Company, Pleasanton, California

[‡] Sabbatical Visitor. Permanent address: Tsing Hua University, Taiwan, NRC

Introduction

The pyrolysis of 1-iodopropane has recently been the subject of a number of VLPP studies.¹⁻³ The mechanism of the decomposition has been discussed in detail by these earlier workers. The molecule decomposes by two channels whose critical reaction thresholds, E_0 , differ by 4.3 kcal mole⁻¹:



In reaction (2), the first step is rate determining.

This chemical system is an attractive one for study by the Variable Encounter Method (VEM) because it affords an opportunity for comparison of our results in the transient region with those obtained in steady state VLPP studies. Moreover, this molecule is the first polar one to be studied by the VEM technique. Information is desirable concerning the dependence of the energy transfer process between molecules and the wall upon this parameter.

In this paper we report the study of the pyrolysis of 1-iodopropane in a VEM system at temperatures from 625 K to 840 K. Unfortunately, the system seems to be not as clean as we had hoped originally.

Experimental

The experimental apparatus was similar to that used in previous VEM studies.^{4,5} Reactor fingers with mean numbers of collisions, m , of 5.9 and 17.5 were used; these were described in ref. 5. Reproducible results (within the precision of the experiments) were obtained after "seasoning" the reactors for several hours. The resulting seasoned surface deteriorated with time when standing under vacuum (unlike our experience with hydrocarbon substrates^{4,5}), or when the system was pumped for long periods, suggesting that the seasoning film itself was either being slowly pyrolyzed or had an appreciable vapor pressure.

Matheson-Coleman 1-iodopropane containing no detectible impurities was thoroughly degassed prior to use. Experiments varied in duration from one minute to several hours, depending on experimental conditions. The reaction was allowed to proceed usually from 1 to 30 per cent completion, and an aliquot was then taken. The products were analyzed by gas chromatography with fid detection. Separation of 1-iodopropane from its hydrocarbon reaction products was made on a 12 inch x 3/16 inch packed column consisting of 15% squalane on 45-60 mesh Chromosorb P. The reaction products were separated on a 6.5 foot x 3/16 inch column having the same packing.

In order to check for possible secondary reactions involving the radicals produced by reaction (2), a series of experiments was performed with trans-2-betene added as a getter. The ratio of iodopropane to butene was varied between 0.1 to 0.2.

Results and Calculations

Experimentally determined values of the mean probability of reaction per collision, $\bar{P}_c(m)$, by channel (1) are plotted versus temperature in Fig. 1 for the two reactors. These results are similar in behavior to those obtained in previous VEM studies. However, the rate data display somewhat more scatter (Fig. 2) than those obtained in previous studies and showed greater sensitivity to the seasoning procedure and history. Systematic fluctuations of $\sim 50\%$ occur. Moreover, small amounts of C_2 impurities, presumably arising from the decomposition of the seasoning film, were always present and made it unattractive to expend much effort on the measurement of reaction rates for reaction (2) which produces ethylene. Decomposition by this latter channel is accompanied by the production of radicals and traces of propane were observed in some of the high temperature experiments. Some differences (10-30%) in the apparent rate constants for propylene formation were observed between the experiments with butene getter and those using neat 1-iodopropane (Fig. 2). However, both negative and positive deviations occur and reflect mainly data scatter.

We note that earlier experimental data obtained in VLPP studies of 1-iodopropane also indicates some apparent experimental complications.^{2,3} In that work, apart from scatter in at least some of the data, comparable to that shown here, the observed k_1/k_2 ratio decreased (rather than the expected constancy) as the number of collisions in the reactor increased; also, rates in a 2140-collision reactor showed a larger temperature dependence than did a 19,950-collision reactor. Additionally, reaction (2), unlike reaction (1), was found to show no dependence on the pressure of added inert bath gas. Not much weight was placed on these complications by the earlier workers. We believe that although experimental complications do affect the quantitative precision of the results, that the general conclusions derived below and based on reaction (1) are

substantially valid.

The RRKM models employed for reactions (1) and (2) were based on those reported by Gaynor et al. (2). If molecules were effectively in the second order region, the only parameter of critical importance would be E_0 , the critical energy for reaction, together with the molecular parameters. However, 1-iodopropane only approaches the second order region under the conditions used in this study. The fraction of molecules activated above E_0 which react between collisions is $\sim 14\%$ at 691 K and increases to $\sim 19\%$ at 804 K. The molecular and activated complex parameters and vibrational frequencies used are listed in Table I. We have altered Gaynor's frequency factor and reaction path degeneracy for reaction (1), taking $\log A_\infty$ to be 13.0 and L^\ddagger to be 1.5. 1-Iodopropane has roughly equal probabilities of being in the trans and gauche forms (6,7); these have reaction path degeneracies for reaction (1) of 2 and 1, respectively. Hence, an overall reaction path degeneracy of 1.5 seems to us more realistic than the reported value of 0.33. Calculated values of the microscopic rate constants k_E are plotted versus energy in Fig. 3.

The computer simulation of the encounter process has been described elsewhere.⁴ Four models have been used to characterize the probability, p_{ij} , of a down-transition by the molecule, from energy E_j to energy E_i :^{5,8}

Model FE (flat exponential, i.e., $\langle \Delta E \rangle$ independent of E_j):

$$p_{ij} = A_1 \exp(-\Delta E / \langle \Delta E \rangle); \quad \Delta E = E_j - E_i$$

Model FG (flat gaussian, i.e., ΔE_{mp} independent of E_j):

$$p_{ij} = A_2 \exp(-(\Delta E - \Delta E_{mp})^2 / 2\sigma^2);$$

Model EB (exponentially weighted Boltzmann):

$$p_{ij} = A_3 N(E_i) \exp(E_i / RT) \exp(-\Delta E / \langle \Delta E \rangle);$$

Model GB (gaussian weighted Boltzmann):

$$p_{ij} = A_4 N(E_i) \exp(-E_i / RT) \exp(-(\Delta E - \Delta E_m)^2 / 2\sigma^2);$$

The A's are normalization constants, $N(E_i)$ is the density of quantum states at energy E_i , and $\langle \Delta E \rangle$, ΔE_{mp} and σ are parameters of the models. In this calculation, σ was given the value of $0.7 \Delta E_{mp}$. For Models FE and FG, a few transitions can occur in principle to energy levels $E_i < 0$, and these were simply treated as elastic. Alternative treatments were explored and do not affect present conclusions. Of these models, FE has been shown to be inappropriate for surfaces that behave as near-strong colliders, as is the case here at the lower temperatures; results for this model will not be shown. Truncation of the transition probabilities for ΔE greater than $18,000 \text{ cm}^{-1}$ (i.e. $p_{ij} = 0$ for $\Delta E \geq 18,000$) was employed for most of the computations but gave results essentially identical to those obtained in calculations where no truncation was employed.

Representative parameters of each of the models which produce a fit to the 5.9 reactor data are listed in Table 2, and the calculated probabilities of reactions per collision in a given reactor, $\bar{P}_c(m)$, are listed in Table 2 and presented in Fig. 1. Also shown is a calculated curve for an ideal strong collider. The parameters derived from the $m = 5.9$ fit were then applied to the calculation of curves for case $m = 17$ and comparison of these with experiment is also shown in Fig. 1.

The effective average energy of down-transitions, $\langle \Delta E' \rangle$ for Models EB and GB is energy dependent; therefore, in order to characterize these models, the average energy of down-transitions is presented for the case $E_j = \frac{2}{3} E_0$ (E_0 for reaction (1)).

Discussion

At lower temperatures (Fig. 1), the $P_c(m)$ values for the reactor $m = 17.5$ appear within experimental uncertainty to be indistinguishable from those for a true strong collider and correspond to the steady state, i.e., to $\bar{P}_c(\infty)$. The values for the $m = 5.9$ reactor are a little lower than the strong collider values and show more unequivocally that ^{the} surface behaves as a rather efficient collider but only approaches near-strong behavior at temperatures below 700 K. Models EB and GB, which effectively reproduce strong collider behavior in the limit of very large values of the relevant parameters ($\langle \Delta E \rangle$ and ΔE_{mp}) appear to give a better fit to the data (Fig. 1) at low temperatures than does FG, although the scatter in the data is large and this prohibits a decisive conclusion.

Despite the experimental uncertainties, it is possible to draw some important qualitative conclusions. Once again the trend observed in previous VEM studies of decreasing step size with increasing temperature is borne out. More importantly, in its interaction with the wall, this molecule seems to be a much stronger collider than the hydrocarbon molecules previously studied by VEM in this temperature range.^{4,5,8} We presume that this results from the larger polarity and stronger attractive molecular potential of the molecule.

We may make a limited comparison of our results with those reported in VLPP studies of 1-iodopropane.^{1,3} The highest temperature at which we were able to obtain reliable data (840 K) is a little lower than the lowest temperature (850 K) at which the VLPP results were reported. But, while Gaynor, et al. report that the rates of reactions (1) and (2) are about equal, our calculated steady state k_2/k_1 ratio (at 805 K) based on the preferred energy transfer models is much lower, 0.29 or less, as shown in Table 3. In fact, based on our experiments, 0.29 is also a maximum in our observed experimental ratios. We are uncertain as to the origin and significance of this difference.

Table I. Molecular Parameters for RRKM Calculations^a

	1-iodopropane	Complex (1)	Complex (2)
frequencies/cm ⁻¹			
	3100(7)	3100(6)	3100(7)
	1450(5)	2200	1450(5)
	1150(6)	1450(4)	1150(4)
	1000(2)	1300	1000(2)
	700	1150(7)	700(2)
	600	1000	400
	425(2)	700(2)	200
	300	300	130(3)
	200	219(3)	70
	70		
$(I_A I_B I_C)^+ / (I_A I_B I_C)$	--	2.83	3.09
L^+	--	1.5	1.0

	$E_0/\text{kcal mole}^{-1}$	$E_{a\infty}/\text{kcal mole}^{-1}$	$\log_{10}(A_{\infty}/\text{s}^{-1})$
Reaction (1)	48.5	49.5	13.0
Reaction (2)	52.8	55.7	15.5

a) Based on refs. 2 and 3, except as in text

Table 2. Calculated Values of $\bar{P}_c(m)$ and Parameters of the Energy Transfer Models

T/K	Model	$\langle \Delta E \rangle$ or ΔE_{mp}	$\langle \Delta E' \rangle$	$\bar{P}_c(5.9)^a$	$P_c(17.5)$
690	FG	3740	4136	5.69-8 ^c	1.19-7
	EB	1370	4113 ^b		1.27-7
	GB	2165	3884 ^b		1.27-7
805	FG	2730	3012	2.62-6	6.38-6
	EB	1335	2664 ^b		7.07-6
	GB	1895	2888 ^b		7.08-6

a) Models fitted to the experimental values for $m = 5.9$.

b) This quantity is energy-level dependent and was calculated for a representative value of $E = \frac{2}{3} E_0$; the quantity increases with increase of E .

c) Signifies 5.69×10^{-8}

Table 3. Calculated Values of k_2/k_1


<u>T/K</u>	<u>Model FE(m = 17.5)</u>	<u>Model EB(m = 17.5)</u>
691	0.14	0.17
805	0.21	0.29

Bibliography


1. K. D. King, D. M. Golden, G. N. Spokes and S. W. Benson, Int. J. Chem. Kinet., 3, 411 (1971).
2. B. J. Gaynor, R. G. Gilbert and K. D. King, Chem. Phys. Lett., 58, 591 (1978).
3. B. J. Gaynor, Ph.D. Dissertation, University of Sydney, (1980).
4. D. F. Kelley, L. Zalotai and B. S. Rabinovitch, Chem. Phys., 46, 379 (1980).
5. M. C. Flowers, F. C. Wolters, B. D. Barton and B. S. Rabinovitch, Chem. Phys., 47, 189 (1980).
6. Y. Ogawa, S. Imazeki, H. Yamaguchi, H. Matura, I. Harada and T. Shimanouchi, Bull. Chem. Soc. Japan, 51, 743 (1978).
7. K. Tanabe and S. Saeki, J. Mol. Struct., 27, 79 (1975).
8. F. C. Wolters, M. C. Flowers and B. S. Rabinovitch, J. Phys. Chem., 85, xxx (1981).

Figure Captions


- Figure 1. Plot of experimental and calculated values of $P_c(m)$ for reaction (1) vs temperature (K) for 1-iodopropane. Experimental results: \bigcirc , $m = 5.9$; \square , $m = 17.5$. Calculated results: curve 1, strong collider; curve 2, Models EB and GB for $m = 17.5$; curve 3, Model FG for $m = 17.5$; curve 4, calculated and experimental values for $m = 5.9$. Error bars show 95% confidence level.
- Figure 2. Plots of $\ln A/\dot{A}$ vs time for several runs, with and without added butene. The ratio of rate constants in the two cases is given in the figure; \bigcirc , 626 K; \triangle , 694 K; \square , 828 K. Filled symbols refer to butene runs. Time is in minutes at 626 K and 694 K and in seconds at 828 K.
- Figure 3. Plot of calculated values of $\log_{10} (k_E/s^{-1})$ vs E/cm^{-1} for reaction (1), (\bullet) , and reaction (2), (x) .



Erst Wollers et al



For better detail



From Walters et al

INT.

ENERGETIC MATERIALS RESEARCHDISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Assistant Secretary of the Navy (R, E, and S) Attn: Dr. R.E. Reichenbach Room 5E787 Pentagon Washington, DC 20350	1	AFATL Eglin AFB, FL 32542 Attn: Dr. Otto K. Heiney	1
Office of Naval Research Code 473 Arlington, VA 22217 Attn: Dr. R. Miller	10	AFRPL Code PACC Edwards AFB, CA 93523 Attn: Mr. W. C. Andrepont	1
Office of Naval Research Code 200B Arlington, VA 22217 Attn: Dr. J. Enig	1	AFRPL Code CA Edwards AFB, CA 93523 Attn: Dr. R. R. Weiss	1
Office of Naval Research Code 260 Arlington, VA 22217 Attn: Mr. D. Siegel	1	Code AFRPL MKPA Edwards AFB, CA 93523 Attn: Mr. R. Geisler	1
Office of Naval Research Western Office 1030 East Green Street Pasadena, CA 91106 Attn: Dr. T. Hall	1	Code AFRPL MKPA Edwards AFB, CA 93523 Attn: Dr. F. Roberto	1
Office of Naval Research Eastern Central Regional Office 495 Summer Street Boston, MA 02210 Attn: Dr. L. Peebles Dr. A. Wood	2	AFSC Andrews AFB, Code DLFP Washington, DC 20334 Attn: Mr. Richard Smith	1
Office of Naval Research San Francisco Area Office One Hallidie Plaza Suite 601 San Francisco, CA 94102 Attn: Dr. P. A. Miller	1	Air Force Office of Scientific Research Directorate of Chemical & Atmospheric Sciences Bolling Air Force Base Washington, DC 20332	1
Defense Technical Information Center DTIC-DDA-2 Cameron Station Alexandria, VA 22314	12	Air Force Office of Scientific Research Directorate of Aero- space Sciences Bolling Air Force Base Washington, DC 20332 Attn: Dr. L. H. Caveny	1
		Anal-Syn Lab Inc. P.O. Box 547 Paoli, PA 19301 Attn: Dr. V. J. Keenan	1

	<u>No. Copies</u>		<u>No. Copies</u>
Army Ballistic Research Labs Code DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Mr. L. A. Watermeier	1	Hercules Inc. Eglin AFATL/DL DL Eglin AFB, FL 32542 Attn: Dr. Ronald L. Simmons	1
Army Ballistic Research Labs ARRADCOM Code DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Dr. Ingo W. May	1	Hercules Inc. Magna Bacchus Works P.O. Box 98 Magna, UT 84044 Attn: Mr. E. H. DeButts	1
Army Ballistic Research Labs ARRADCOM Code DRDAR-BLT Aberdeen Proving Ground, MD 21005 Attn: Dr. Philip Howe	1	Hercules Inc. Magna Bacchus Works P.O. Box 98 Magna, UT 84044 Attn: Dr. James H. Thacher	1
Army Missile Command Code DRSME-RK Redstone Arsenal, AL 35809 Attn: Dr. R. G. Rhoades Dr. W. W. Wharton	2	HQ US Army Material Development Readiness Command Code DRCDE-DW 5011 Eisenhower Avenue Room 8N42 Alexandria, VA 22333 Attn: Mr. S. R. Matos	1
Atlantic Research Corp. 5390 Cherokee Avenue Alexandria, VA 22314 Attn: Dr. C. B. Henderson	1	Johns Hopkins University APL Chemical Propulsion Information Agency Johns Hopkins Road Laurel, MD 20810 Attn: Mr Theodore M. Gilliland	1
Ballistic Missile Defense Advanced Technology Center P.O. Box 1500 Huntsville, AL 35807 Attn: Dr. David C. Sayles	1	Lawrence Livermore Laboratory University of California Livermore, CA 94550 Attn: Dr. M. Finger	1
Ballistic Research Laboratory USA ARRADCOM DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Dr. A. W. Barrows	1	Lawrence Livermore Laboratory University of California Livermore, CA 94550 Attn: Dr. R. McGuire	1
Hercules Inc. Cumberland Aerospace Division Allegany Ballistics Lab P.O. Box 210 Cumberland, MD 21502 Attn: Dr. Rocco Musso	2	Lockheed Missiles and Space Co. P.O. Box 504 Sunnyvale, CA 94088 Attn: Dr. Jack Linsk Org. 83-10 Bldg. 154	1

	<u>No. Copies</u>		<u>No. Cop</u>
Lockheed Missile & Space Co. 3251 Hanover Street Palo Alto, CA 94304 Attn: Dr. H. P. Marshall Dept. 52-35	1	Naval Research Lab Code 6100 Washington, DC 20375	1
Los Alamos Scientific Lab P.O. Box 1663 Los Alamos, NM 87545 Attn: Dr. R. Rogers, WX-2	1	Naval Sea Systems Command Washington, DC 20362 Attn: Mr. G. Edwards, Code 62R3 Mr. J. Murrin, Code 62R2 Mr. W. Blaine, Code 62R	1
Los Alamos Scientific Lab P.O. Box 1663 Los Alamos, NM 87545 Attn: Dr. B. Craig, M Division	1	Naval Sea Systems Command Washington, DC 20362 Attn: Mr. R. Beauregard SEA 64E	1
Naval Air Systems Command Code 330 Washington, DC 20360 Attn: Mr. R. Heitkotter Mr. R. Brown	1	Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. H. G. Adolph	1
Naval Air Systems Command Code 310 Washington, DC 20360 Attn: Dr. H. Mueller Dr. H. Rosenwasser	1	Naval Surface Weapons Center Code R13 White Oak, Silver Spring, MD 20910 Attn: Dr. R. Bernecker	1
Naval Explosive Ordnance Disposal Facility Indian Head, MD 20640 Attn: Lionel Dickinson Code D	1	Naval Surface Weapons Center Code R10 White Oak, Silver Spring, MD 20910 Attn: Dr. S. J. Jacobs	1
Naval Ordnance Station Code 5034 Indian Head, MD 20640 Attn: Mr. S. Mitchell	1	Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. M. J. Kamlet	1
Naval Ordnance Station Code PM4 Indian Head, MD 20640 Attn: Mr. C. L. Adams	1	Naval Surface Weapons Center Code R04 White Oak, Silver Spring, MD 20910 Attn: Dr. D. J. Pastine	1
Dean of Research Naval Postgraduate School Monterey, CA 93940 Attn: Dr. William Tolles	1	Naval Surface Weapons Center Code R13 White Oak, Silver Spring, MD 20910 Attn: Dr. E. Zimet	1
Naval Research Lab Code 6510 Washington, DC 20375 Attn: Dr. J. Schnur	1		

	<u>No. Copies</u>		<u>No. Copies</u>
Naval Surface Weapons Center Code R101 Indian Head, MD 20640 Attn: Mr. G. L. MacKenzie	1	Naval Weapons Center Code 388 China Lake, CA 93555 Attn: D. R. Derr	1
Naval Surface Weapons Center Code R17 Indian Head, MD 20640 Attn: Dr. H. Haiss	1	Naval Weapons Center Code 388 China Lake, CA 93555 Attn: Dr. R. Reed Jr.	1
Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. K. F. Mueller	1	Naval Weapons Center Code 385 China Lake, CA 93555 Attn: Dr. A. Nielsen	1
Naval Surface Weapons Center Code R16 Indian Head, MD 20640 Attn: Dr. T. D. Austin	1	Naval Weapons Center Code 3858 China Lake, CA 93555 Attn: Mr. E. Martin	1
Naval Surface Weapons Center Code R122 White Oak, Silver Spring, MD 20910 Attn: Mr. L. Roslund	1	Naval Weapons Center China Lake, CA 93555 Attn: Mr. R. McCarten	1
Naval Surface Weapons Center Code R121 White Oak, Silver Spring, MD 20910 Attn: Mr. M. Stosz	1	Naval Weapons Support Center Code 5042 Crane, Indiana 47522 Attn: Dr. B. Douda	1
Naval Weapons Center Code 3853 China Lake, CA 93555 Attn: Dr. R. Atkins	1	Rohm and Haas Company 723-A Arcadia Circle Huntsville, Alabama 35801 Attn: Dr. H. Shuey	1
Naval Weapons Center Code 3205 China Lake, CA 93555 Attn: Dr. L. Smith	1	Strategic Systems Project Office Dept. of the Navy Room 901 Washington, DC 20376 Attn: Dr. J. F. Kincaid	1
Naval Weapons Center Code 3205 China Lake, CA 93555 Attn: Dr. C. Thelen	1	Strategic Systems Project Office Dept. of the Navy Room 1048 Washington, DC 20376 Attn: Mr. E. L. Throckmorton Mr. R. Kinert	2
Naval Weapons Center Code 385 China Lake, CA 93555 Attn: Dr. A. Amster	1	Thiokol Chemical Corp. Brigham City Wasatch Division Brigham City, UT 84302 Attn: Dr. G. Thompson	1

	<u>No. Copies</u>		<u>No. Copies</u>
USA ARRADCOM DRDAR-LCE Dover, NJ 07801 Attn: Dr. R. F. Walker	1	University of California Department of Chemistry 405 Hilgard Avenue Los Angeles, CA 90024 Attn: Professor M. Nicol	1
USA ARRADCOM DRDAR-LCE Dover, NJ 07801 Attn: Dr. N. Slagg	1	University of California Energy Center Mail Code B-010 La Jolla, CA 92093 Attn: Prof. S.S. Penner	1
U.S. Army Research Office Chemistry Division P.O. Box 12211 Research Triangle Park, NC 27709	1		
Washington State University Dept. of Physics Pullman, WA 99163 Attn: Professor G.D. Duval	1	Dr. P. Rentzepis Bell Laboratories Murray Hill N.J. 07971	1
Space Sciences, Inc. 135 West Maple Avenue Monrovia, CA 91016 Attn: Dr. M. Farber	1	University of Southern CA Department of Electrical Engineering University Park Los Angeles, CA 90007 Attn: C. Wittig	1
SRI International 333 Ravenswood Avenue Menlo Park, CA 94025 Attn: Mr. M. Hill	1	MIT Dept. of Chemistry Cambridge, MA 02139 Attn: Prof. John Deutsch	1
Office of Naval Research Code 421 Arlington, VA 22217 Attn: Dr. B. Junker	1		
The Johns Hopkins University Department of Chemistry Baltimore, MD 21218 Attn: Dr. Joyce J. Kaufman	1		
University of California Department of Chemistry Berkeley, CA 94720 Attn: Professor Y.T. Lee	1		
Office of Naval Research Code 472 Arlington, VA 22217 Attn: Dr. G. Neece	1		